

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 732 403 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
18.09.1996 Bulletin 1996/38

(51) Int. Cl.⁶: **C12N 15/81**
// C07K14/02

(21) Application number: **96200286.1**

(22) Date of filing: **06.01.1984**

(84) Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

(30) Priority: **22.02.1983 US 468589**

(62) Application number of the earlier application in
accordance with Art. 76 EPC: **91114001.0**

(71) Applicant: **CHIRON CORPORATION**
Emeryville, California 94608 (US)

(72) Inventors:
• **Burke, Rae L.**
San Francisco, California 94114-2114 (US)
• **Rosenberg, Steven**
Oakland, California 94602 (US)

• **Tekamp-Olson, Patricia**
San Francisco, California (US)
• **Valenzuela, Pablo D.T.**
San Francisco, California (US)

(74) Representative: **Hallybone, Huw George et al**
CARPMAELS AND RANSFORD
43 Bloomsbury Square
London WC1A 2RA (GB)

Remarks:

This application was filed on 08 - 02 - 1996 as a
divisional application to the application mentioned
under INID code 62.

(54) **Yeast expression systems with vectors having GAPDH promoters, and synthesis of HBsAg**

(57) Yeast cells containing DNA plasmids having
foreign DNA, wherein foreign DNA is expressed, are
described. For example, DNA coding for hepatitis B and
its virus surface antigen (HBsAg) is ligated to a yeast
plasmid to yield a product that is used to transform
yeast cells. The plasmids of this invention have GAPDH
promoters, and are capable of replicating in either a
yeast cell or a bacterial cell.

EP 0 732 403 A1

Description

BACKGROUND OF THE INVENTION

5 For maximal expression of foreign genes in microbial systems it is usually advantageous to employ homologous regulatory elements within the expression vector. Efficiency of expression (product formation) is believed to be a function of and proportional to the strength of the promoter employed. In addition, regulation of gene expression by nutritional factors under the control of the experimenter offers a further useful manipulatory tool. The glycolytic enzyme genes of yeast, e.g., those coding for glyceraldehyde-3-phosphate dehydrogenase (GAPDH) and pyruvate kinase (PyK), possess the above useful properties, i.e., high levels of expression (and thus by inference very efficient promoters) and susceptibility to regulation by components of the growth medium. For example, GAPDH can comprise as much as 5% of the dry weight of commercial baker's yeast (Krebs, E.G., *J. Biol. Chem.* (1953) 200:471). Furthermore, these enzymes are also highly inducible. For example, when yeast cultures are shifted from growth on acetate to glucose, the activity of GAPDH increased up to 200-fold in proportion to the concentration of the sugar in the medium (Maitra, P.K. and Lobo, Z., *J. Biol. Chem.* (1971) 246:475). These results suggest that the transcriptional machinery of these genes is highly regulated, perhaps by the participation of DNA sequences present in the 5' non-coding flanking region of the genes.

This invention relates to the isolation, structure and the successful use in yeast expression plasmids of DNA fragments corresponding to the 5' non-coding regions of the regulatable yeast genes GAPDH and PyK. These fragments which contain DNA sequences with strong transcription-promoting activity are called "promoters". They are ideal components of DNA vectors for commercial production of large quantities of protein coded by foreign genes under their transcriptional control.

In addition, this invention encompasses yeast expression plasmids further comprising an appropriate terminator to a form a "cassette" of promoter-foreign gene-terminator. The presence of the terminator increases expression of the foreign DNA.

An early attempt to express foreign DNA in yeast failed (Beggs, J.D. *et al.*, *Nature* (1980) 283:285). In this report, the hemoglobin DNA (inserted with its own promoter) was transcribed but the RNA was not spliced. A variety of explanations for this result are possible, e.g., an incorrect location for the initiation of transcription and/or the poor ability of yeast cells to carry out splicing of intervening sequences (introns).

30 Three GAPDH genes of yeast have been cloned (Holland, M.J. *et al.*, *Basic Life Science* (1981) 19:291), but their promoters have not been used for constructing expression systems in yeast by recombinant DNA methods. The PyK gene has also been cloned, but by genetic complementation only (no structural studies performed) (Kawasaki, G. and Fraenel, D.G., *Biochem. Biophys. Res. Comm.* (1982) 108:1107). Other yeast promoters, e.g., that of alcohol dehydrogenase I (Valenzuela, P. *et al.*, *Nature* (1982) 298:347 and Hitzeman, R.A. *et al.*, *Nature* (1981) 293:717) and phosphoglycerate kinase (Tuite, M.F. *et al.*, *EMBO J.* (1982) 1:603 and Hitzeman, R.A. *et al.*, *Science* (1983) 219:620) have been linked to foreign genes to produce yeast expression but no terminators were used. The present invention provides new promoters for yeast expression systems and combines the advantages of highly expressive promoters with the enhanced expression found with appropriately ligated terminators.

A published European Patent Application (072 318) disclosed the construction of a yeast expression vector which, upon induction, expressed hepatitis B virus surface antigen (HBsAg) S-protein under control of the yeast alcohol dehydrogenase I (ADHI) promoter.

BRIEF DESCRIPTION OF THE INVENTION

45 This invention relates to a yeast expression vector comprising a segment of foreign DNA, e.g., that coding for hepatitis B virus (HBV) surface antigen (HBsAg), under transcriptional control of a yeast GAPDH promoter. Terminators may also be appropriately attached. The expression vector typically has a yeast replication origin and is capable of replicating in either type of cell. The expression vector, when used to transform yeast cells, will yield substantial amounts of the protein coded by the segment of foreign DNA.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1: Isolation and tailoring of a GAPDH promoter fragment.

Fig. 2: DNA sequences of the GAPDH promoter fragment.

55 Fig. 3: Construction of a yeast expression plasmid containing the GAPDH promoter.

DETAILED DESCRIPTION OF THE INVENTION

In principle, yeast expression plasmids have particular advantages, including the following. Yeast can be grown in large-scale culture for commercial production by processes well-known in the art. In contrast, bacteria in large-scale culture are subject to the frequent problem of "phage-out". Yeast also appears to have much the same ability as mammalian cells to add carbohydrate groups to newly synthesized proteins, a capacity that bacteria do not have. Now that cDNA sequences are readily obtainable, the problem of expressing genes having introns is easily avoided.

The vectors of the present invention encompass promoters of unusually high efficiency. A promoter is defined herein as a DNA segment capable of functioning to initiate transcription of an adjoining DNA segment. Transcription is the synthesis of RNA (herein termed messenger RNA or mRNA), complementary to one strand of the DNA adjoining the promoter region. In eukaryotes, messenger RNA synthesis is catalyzed by an enzyme termed RNA polymerase II. The minimum essential elements of promoter function are the following: To provide a starting point for the initiation of transcription and to provide a binding site for RNA polymerase II near the start site permitting selection of the proper strand of DNA as a template for messenger RNA synthesis. In addition, a eukaryotic promoter functions to regulate the relative efficiency of transcription of coding segments under its control. An active promoter is one which elicits synthesis of relatively large amounts of mRNA complementary to a strand of the adjacent DNA coding segment.

The structural correlates of promoter function have not been clearly established. A promoter segment usually can be identified in nature as a region lying adjacent to the 5' end of a given structural gene. (References to the 5' and 3' ends of a gene will be understood to indicate the corresponding respective ends of mRNA transcribed therefrom, and these, in turn, will be understood to correlate with the NH₂- and -COOH termini of the encoded protein, respectively.) Comparisons of the nucleotide sequences of promoters for various genes from various species have revealed only a few short regions of nucleotide sequence similarity in common among them. Most notable of these is the "TATA Box," a segment of about 5 to 10 nucleotides located generally about 70 to 230 nucleotides upstream from the site of transcription initiation, having a sequence generally resembling TATAA. For review of structural comparisons see Breathnach, R. and Chambon, P., Ann. Rev. of Biochem. (1981) 50:349. The TATA Box is believed to function in initiation of transcription.

The foreign gene will be free or substantially free of codons from the normal structural gene associated with the promoter. Usually, the foreign gene will be joined to a non-coding 3'-end of the regulatory region encompassing the promoter, so as to be free of the amino acids at the N-terminus of endogenous gene naturally associated with the regulatory region. That is, fewer than about 3 codons (9 nucleotides) will be retained with the regulatory region when joined to the foreign gene.

The presence of the terminator sequence at the 3' end of the coding segment enhances expression. The effect is generally similar to the addition of rho factor to prokaryotic transcription systems, wherein the rate of the release of RNA polymerase is enhanced to produce an increase in the rate of reinitiation of transcription. It will be understood that, while the terminator sequences are not required for detectable expression of foreign DNA segments, it is preferable to appropriately link them to enhance expression. The terminator region may be naturally associated with the same or different structural gene as the promoter region.

The most appropriate DNA vector for the GAPDH construction of this invention is a shuttle vector. These vectors can "shuttle" between a bacterial strain, such as E. coli, and yeast, since they have a bacterial origin of replication and a yeast origin of replication, see, e.g., Ammerer, G. et al., Recombinant DNA, Proc. Third Cleveland Symposium Macromolecules (Walton, A.G., ed.), p. 185, Elsevier, Amsterdam (1981). A typical bacterial origin of replication is derived from, e.g., pBR322. The most useful yeast origin of replication is found in the extrachromosomal genetic element known as the 2 micron circle. In laboratory strains the 2 micron plasmid DNA is found in approximately 50 copies per cell and is stably maintained. For a review, see, for example, Curr. Topics Micro. Imm. (1982) 96:119. This yeast plasmid has also been sequenced (Hartley, J.L. et al., Nature (1980) 286:860).

Representative samples of the plasmids and host cells used in the constructions of this invention have been placed on deposit with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland. Plasmid pPyK 9.1.1 and yeast cell transformants 2150-2-3/pHBS-56 GAP347/33 and 2150-2-3/pHBS56PyK were placed on deposit on February 18, 1983 and have received ATCC Accession numbers 40061, 20665 and 20666, respectively.

In the Examples that follow, many of the techniques, reactions and separation procedures are already well-known in the art. All enzymes, unless otherwise stated, are available from one or more commercial sources, such as New England Biolabs, Beverly, Massachusetts; Collaborative Research, Waltham, Massachusetts; Miles Laboratories, Elkhart, Indiana; Boehringer Biochemicals, Inc., Indianapolis, Indiana and Bethesda Research Laboratories, Rockville, Maryland. Buffers and reaction conditions for restriction enzyme digestion were used according to recommendations supplied by the manufacturer for each enzyme, unless otherwise indicated. Standard methodology for other enzyme reactions, gel electrophoresis separations and E. coli transformation may be found in Methods in Enzymology, (1979) 68. Transformation of yeast protoplasts can be carried out essentially as described by Beggs, Nature (1978) 275:104.

E. coli strains useful for transformation include X1776; K12 strain 294 (ATCC No. 31446); RR1 and HB101. Yeast strains XV610-8c having the genotype (a ade2 ade6 leu2 lys1 trp1 can1) and GM-3C-2, genotype: (Leu2 Trp1 His4

CYC1-1CYP3-1) (Faye, G. *et al.*, *Proc. Natl. Acad. Sci.* (1981) 78:2258) can be typically used for yeast transformations. It would be understood, however, that virtually any strain of yeast is useful for transformation. Bacteria can be grown and selected according to procedures described by Miller, J.H., *Experiments in Molecular Genetics*, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York (1972). Yeast can be grown on the following media: YEPD containing 1% (w/v) yeast extract, 2% (w/v) peptone and (w/v) glucose; and, in the case of plating medium, 3% (w/v) agar. YNB plus CAA contains 6.7 grams of yeast nitrogen base (Difco Laboratories, Minneapolis, Minnesota), 10mg of adenine, 10mg of uracil, 5g casamino acids (CAA) (Difco), 20g glucose; and, in the case of plating media, 30g agar per liter. Selection for tryptophan prototrophy can be made on plates containing 6.7g yeast nitrogen base (lacking amino acids), supplemented for all growth requirements of the strain to be transformed except tryptophan.

EXAMPLE 1

Cloning of the yeast glyceraldehyde-3-phosphate dehydrogenase (GAPDH) gene.

A complementary DNA (cDNA) containing the yeast GAPDH coding sequences was prepared in the following manner:

PolyA⁺ RNA was isolated from yeast strain A364A. Double-stranded cDNA was synthesized using AMV reverse transcriptase and *E. coli* DNA polymerase I. Poly-dC-tails were added to the double-stranded cDNA molecule using deoxynucleotide terminal transferase. Poly-dC-tailed cDNA was annealed to poly-dG-tailed pBR322 and used to transform *E. coli* HB101. One thousand transformants were screened by colony hybridization to labeled PolyA⁺ RNA, and a subset further examined by restriction endonuclease mapping, and DNA sequencing. Three clones containing GAPDH sequences were isolated from the pool. One clone (pcGAP-9) contained an insert of about 1200 base pairs (bp) and was used for further work.

A yeast gene library was prepared by inserting fragments obtained after partial digestion of total yeast DNA with restriction endonuclease *Sau*3A into lambda phage Charon 28, according to Blattner, F.R. *et al.*, *Science* (1977) 196:161-169. Several fragments containing yeast GAPDH coding sequences were isolated by screening the phage library with labeled DNA from pcGAP-9. The yeast GAPDH gene of one of these clones was subcloned in pBR322 as a 2.1kb *Hind*III fragment (pGAP-1, see Fig. 1) or as a 3.5kb *Bam*HI fragment (pGAP-2). The GAPDH promoting-active fragments were isolated from these clones. The *Hind*III-*Hha*I fragment of about 800bp was ligated to the *Hha*I-*Hind*III fragment of about 350bp. The resulting 1061bp *Hind*III fragment was isolated by gel electrophoresis and cloned in pBR322, (pGAP-347), and the sequence determined (see Fig. 2).

EXAMPLE 2

Construction of yeast vectors containing the GAPDH promoter, active in the expression of HBsAg.

A plasmid vector (pHBS-56GAP347/33), for the expression of HBV surface antigen in yeast, using the GAPDH promoter fragment was constructed as depicted in Fig. 3.

Total digestion of pGAP-347 with *Sph*I followed by partial digestion with *Hind*III yielded an approximately 1700bp *Sph*I-*Hind*III fragment having about 1060bp of GAPDH promoter and about 530bp of pBR322. The 1700bp *Sph*I-*Hind*III GAPDH promoter fragment was ligated with the 840bp *Hind*III-*Hind*III fragment (containing the HBsAg coding region, 26 bases of 5' non-coding region and 128bp of 3' non-coding region, obtained from pHBS-56) and then with the 350bp *Hind*III-*Sph*I fragment containing the ADH-1 termination region (isolated from pHBS-56). The 2900bp *Sph*I fragment (cassette) was isolated and cloned in pHBS-56 previously digested with *Sph*I. The plasmid pHBS-56 (ATCC Accession No. 40047) has been described in a co-pending application (EPA No. 82.401473.2 published as no. 72318, of Regents of the University of California, herein incorporated by reference) and contains the entire 2 micron plasmid, in addition to a region with the yeast *leu2* gene and the amp resistance locus of pBR322. The resulting plasmid (pHBS-56GAP347/33) in which the promoter, gene and termination regions were in the proper orientations was isolated and used to transform yeast strain AB102 (*MATa*, *pep* 4-3, *leu* 2-3 *leu*2-112, *ura* 3-52, *his* 4-580, *cir*^o) or strain 2150-2-3 (*MATa*, *ade*1, *leu*2-04, *cir*^o). Strain AB102 is derived from SF657-9c by curing of 2 micron plasmids. Strain 2150-2-3 is from the collection of Dr. Leland Hartwell at the University of Washington.

EXAMPLE 3

Synthesis of HBsAg in yeast under GAPDH promoter control (plasmid pHBS-56GAP347/33).

One hundred ml cultures of strain AB102 containing plasmid pHBS56-347/33 were grown to optical density at 650nm of 1. Cell-free lysates were prepared by agitation with glass beads and removal of cell debris by centrifugation. HBsAg was measured by the Abbott Ausriall radioimmunoassay and protein concentration was determined by the

Coomassie blue binding method. The results are shown in Table 1. They indicate that the GAPDH promoter is about 5 times more effective than the ADH-1 promoter for protein product expression in yeast.

Table 1

Synthesis of HBsAg in yeast			
(a) control front pHBS-56 (ADH-I promoter)			
Exp#	sAg (µg/ml)	protein (mg/ml)	Spec. Activity (µgsAg/mg protein)
1	8.8	18	0.49
2	14	25	0.56
3	12.4	20	0.62
(b) from pHBS-56GAP347/33 (GAPDH promoter)			
Exp#	sAg (µg/ml)	protein (mg/ml)	Spec. Activity (µgsAg/mg protein)
1	36	14	2.6
2	35	12	2.9
3	37	12.5	3.0

Similar results were obtained by substituting yeast strain 2150-2-3 for yeast strain AB102 and repeating Example 3.

Claims

1. A yeast expression vector comprising a segment of foreign DNA under transcriptional control of a yeast glyceraldehyde-3-phosphate dehydrogenase promoter, said segment being in the correct orientation for transcription and having three or fewer codons from yeast glyceraldehyde-3-phosphate dehydrogenase at the 5'-end of said foreign DNA, wherein said foreign DNA encodes hepatitis B surface antigen or a portion thereof.
2. A yeast expression vector of claim 1, further comprising a terminator attached to the 3' end of the segment of foreign DNA.
3. A yeast expression vector of any of claims 1 and 2, further comprising a yeast cell replication origin, and being capable of replicating in a yeast cell.
4. A yeast expression vector of any of claims 1, 2 and 3, further comprising a bacterial cell replication origin, and being capable of replicating in a bacterial cell.
5. A yeast expression vector according to claim 2, wherein said terminator comprises the yeast alcohol dehydrogenase I terminator.
6. A yeast expression vector according to claim 2, wherein said terminator comprises the yeast GAPDH terminator.
7. A yeast expression vector according to claim 2, wherein said terminator comprises the yeast PyK terminator.
8. A yeast expression vector of claim 1, further comprising yeast two micron plasmid DNA or a portion thereof.
9. A method of expressing a DNA coding segment in yeast, comprising the steps of:
 - (a) inserting in a yeast expression vector a coding segment which encodes hepatitis B surface antigen or a portion thereof, said vector comprising a DNA segment derived from a yeast glyceraldehyde-3-phosphate dehydrogenase promoter having three or fewer codons from the 5'-end of yeast glyceraldehyde-3-phosphate dehydrogenase, said promoter being adjacent to the 5' end of the inserted DNA coding segment and so oriented that transcription initiated within said promoter includes the coding segment, thereby providing a coding segment expression vector, and
 - (b) transforming yeast cells with the coding segment expression vector.

10. A method according to claim 9, wherein said yeast expression vector further comprises a terminator attached to the 3' end of the inserted DNA coding segment.

11. A method according to claim 10, wherein said terminator comprises the yeast alcohol dehydrogenase I terminator.

12. A method according to claim 10, wherein said terminator comprises the yeast GAPDH terminator.

13. A method according to claim 10, wherein said terminator comprises the yeast PyK terminator.

14. A method according to any of claims 10, 11, 12, and 13, wherein said yeast expression vector further comprises a yeast cell replication origin and is capable of replicating in a yeast cell.

15. A method according to any of claims 10, 11, 12, and 13, wherein said yeast expression vector further comprises a bacterial cell replication origin and is capable of replicating in a bacterial cell.

16. A method according to any of claims 10, 11, 12, and 13, wherein said yeast expression vector further comprises a yeast cell replication origin and a bacterial cell replication origin and is capable of replicating in yeast cells or bacterial cells.

17. A method according to any of claims 10, 11, 12, and 13, wherein said yeast expression vector further comprises yeast two micron plasmid DNA or a portion thereof.




FIG. 1.

DNA 347

10	20	30	40	50	60
AAGCTTACCA	GTTCCTCACAC	GGAACACCAC	TAAATGGACAC	AAATTCGAAA	TACTTTGACC
70	80	90	100	110	120
CTATTTTCGA	GGACCTTGTC	ACCTTGAGCC	CAAGAGAGCC	AAGATTTAAA	TTTTCCTATG
130	140	150	160	170	180
ACTTGAIGCA	AATTCCTAAA	GCTAATAACA	TGCAAGACAC	GTACGGTCAA	GAAGACATAT
190	200	210	220	230	240
TTGACCTCTT	AACTGGTTCA	GACGCGACIG	CCTCATCAGT	AAGACCCGTT	GAAAAGAAGT
250	260	270	280	290	300
TACCTGAAAA	AAACGAATAT	ATACTAGCGT	TGAATGTTAG	CGTCAACAAC	AAGAAGTTTA
310	320	330	340	350	360
ATGACGCGGA	GGCCAAGGCA	AAAAGATICC	TTGATTACGT	AAGGGAGTTA	GAATCATTTT
370	380	390	400	410	420
GAATAAAAAA	CACGCTTTTT	CAGTTCGAGT	TTATCATTAT	CAATACTGCC	ATTTCAAAGA
430	440	450	460	470	480
ATACGTAAAT	AATTAATAGT	AGTGATTTTC	CTAACTTTAT	TTAGTCAAAA	ATTAGCCTTT
490	500	510	520	530	540
TAATTCCTCT	GTAACCCGTA	CATGCCCAAA	ATAGGGGGCG	GGTTACACAG	AATATATAAC
550	560	570	580	590	600
ATCGTAGGTG	TCTGGGTGAA	CAGTTTATCC	CTGGCATCCA	CTAAATATAA	TGGAGCTCGC
610	620	630	640	650	660
TTTTAAGCTG	GCATCCAGAA	AAAAAAGAA	TCCCAGCACC	AAAATATTGT	TTTCTTCACC
670	680	690	700	710	720
AACCATCAGT	TCATAGGTCC	ATTCTCTTAG	CGCAACTACA	GAGAACAGGG	GCACAAACAG
730	740	750	760	770	780
GCAAAAACG	GGCACAACCT	CAATGGAGTG	ATGCAACCTG	CCTGGAGTAA	ATGATGACAC
790	800	810	820	830	840
AAGGCAATTG	ACCCACGCAT	GTATCTATCT	CATTTTCTTA	CACCTTCTAT	TACCTTCTGC
850	860	870	880	890	900
TCTCTCTGAT	TTGGA AAAAG	CTGAAAAAAA	AGGTTGAAAC	CAGTTCCCTG	AAATTATTC
910	920	930	940	950	960
CCTACTTGAC	TAATAAGTAT	ATAAAGACGG	TAGGTATTGA	TTGTAATTCT	GTAAATCTAT
970	980	990	1000	1010	1020
TTCITAAACT	TCTTAAATTC	TACTTTTATA	GTTAGTCTTT	TTTTTAGTTT	TAAACACCA
1030	1040	1050	1060		
ASAACTTAGT	TTCGAATAAA	CACACATAAA	CAACCAAGCT	T	

FIG. 2




FIG. 3.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 96 20 0286

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Y	EP-A-0 060 057 (GENENTECH, INC. & BOARD OF REGENTS UNIVERSITY OF WASHINGTON) 15 September 1982 * page 25, line 14 - line 22 *	1-17	C12N15/81 //C07K14/02
Y	EP-A-0 072 318 (BOARDS OF REGENTS UNIVERSITIES OF CALIFORNIA & WASHINGTON) 16 February 1983 * page 47, line 10 - line 12; claims 1-6; example 2 *	1-17	
A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 80, no. 1, January 1983, WASHINGTON US, pages 1-5, XP002007971 A.MIYANOHARA ET AL.: "Expression of hepatitis B surface antigen in yeast" * the whole document *	1-17	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C12N
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 9 July 1996	Examiner Cupido, M
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.82 (P04C01)